

Changes in Vegetation and Soil Properties - A Case Study of a Quartz Quarry

Vladimir ILINKIN¹ (✉)

Violeta DIMITROVA²

Summary

The paper presents results of an investigation on Technosols. The soils are located in “Podgumer” quartz quarry, village of Podgumer, district of Sofia, Bulgaria. Samples were collected upon deposit, 5 and 9 years after deposit. A phytocenological description was made 5 years after deposit and 9 years after deposit. Soil samples were taken from forty-five soil pits at a depth of 0-20 cm. The vegetation in the area is related to the rhizome - leguminous stage. There are statistically significant differences (a downward trend) in the bulk density of the samples collected 5 and 9 years after deposit. There is a statistically significant increase in total porosity after the establishment of vegetation. The presence of vegetation also results in statistically significant changes in the chemical characteristics (pH, SOM, TKN, P_2O_5 , K_2O). The establishment of vegetation and the increase in plant diversity result in the increase in TKN, P_2O_5 , K_2O , where existing changes have been described using reliable linear regressions.

Key words

technogenic soils, vegetation, soil physical and chemical changes, regression, quartz quarry

¹ Institute of Soil Science, Agrotechnologies and Plant Protection “Nikola Poushkarov”, 7, Shosse Bankya St., Sofia 1331, Bulgaria

² University of Forestry, 10, Kliment Ohridski blvd., Sofia 1756, Bulgaria

✉ Corresponding author: ilinkin@mail.bg

Received: September 5, 2020 | Accepted: November 3, 2020

INTRODUCTION

Human activities have brought about significant changes in the Earth's surface (more than 30% of the Earth's surface), with more than two thirds of these areas being located in Europe (Primack et al., 2012; Chenota et al., 2017). In the early 1980s under the pressure of local communities mainly in developed countries, the issue of landscape quality after mining activities became a matter of concern in projects (Schulz, 1996; Corry et al., 2010). The large number of disused quarries or of those being closed down in numerous countries poses a challenge for the restoration of these severely disturbed habitats (Bradshaw, 1997; Darwish et al., 2011; Mouflis et al., 2008). Different ecosystems develop as a result of the interactions among organisms and their responses to geological, topographic, hydrological and weather conditions. Any disturbance of these conditions can adversely impact the habitats (Nieman and Merkin, 1995). Mining activities are one of the oldest causes for changes in the Earth's surface (Larson et al., 2006; Hristova, 2013). Unlike underground mining where the impact is more localized (Bradshaw, 1997), open-pit mining has a greater environmental impact. The mining industry drastically changes the physical and biological characteristics of the mining areas by destroying flora and fauna, changing soil characteristics and limiting soil microbiocenoses. Thus, abandoned mine lands prove difficult to be revitalized and reused (Corbett et al., 1996). Habitats may be adversely impacted by ecological severance (Nieman and Merkin, 1995), and ecosystem degradation in most post-mining areas is mainly caused by vegetation destruction (Singh et al., 2004; De Deyn et al., 2003). The environmental disturbances that are the most difficult to reclaim are the ones caused by quartz mining. This is mostly due to their specific characteristics and to the light colour of the terrain, which greatly reduces landscape quality and makes it visible even from a very far distance (Pietro et al., 2006; Prieto et al., 2005). Ensuring the sustainable development of ecosystems requires a study of all ecosystem components (Bradshaw and Hüttel, 2001), particularly the soil (Heneghan et al., 2008), being an important, and non-renewable resource (Van-Camp et al., 2004). The diversity of plant species is limited by soil characteristics (Bogdanov, 2013; Bogdanov, 2014a; Hristov, 2020), a vital aspect of which are the physical characteristics of substrates (Romana, 1985). The soil structure in mining areas can easily be damaged. A reduction in porosity results in poor soil aeration, poor drainage and increased resistance to root penetration. This subsequently suppresses plant growth (Bowen et al., 2005; Geissen et al., 2013; Stahl et al., 2002). A positive correlation exists between the availability of nutrients such as C, N, P (total quantities) and the diversity of species. There is an increased accumulation of organic matter when the species are more diverse (Berendse et al., 1998). Microclimate change also affects the decomposition of organic matter (Torreta and Takeda, 1999; Deng et al., 2009). However, the content of organic C, N, P, K doesn't change up to 7 years after mining activities have ceased. The composition of the self-established vegetation includes mainly grass species (Jian-Gang Yuan et al. 2006). The active soil acidity on the territory of quarries is from highly acidic to acidic (3.76- 6.21). There is low nutrient availability (organic C, total N, and total P) (Zhuang, 1997), but it is higher in the surface 0-5 cm layer than in the 5-10 cm layer, where the values vary over time (Sourkova et al., 2005; Pietrzykowski and Krzaklewski, 2007).

MATERIALS AND METHODS

The object. Technosols (WRB, 2014) occurring on the territory of "Podgumer" quartz sand quarry, located next to Podgumer village, district Sofia, Bulgaria. The object is in the lower forest vegetation zone (0 - 600 m a. s. l., Zahariev, 1979).

Methods of study. Forty-five soil pits at a depth of 0-20 cm (in the year of deposit: sample plot 1 – fifteen pits; 5 years after deposit: sample plot 2 – fifteen pits; and 9 years after deposit: sample plot 3 – fifteen pits) were dug at representative plots. A systematic sampling design was used (Petersen & Calvin, 1996). The following soil characteristics were analyzed by using the respective methods:

- Bulk density (BD, g.cm⁻³), according to the DIN ISO 11272:1998, 2001;
- Total porosity (TP, %) according to the Lorraine & Flint (2002);
- Soil texture (Sand - 2 mm - 63 µm, %; Silt - 63 µm - 2 µm, %; Clay < 2 µm, %) using the sedimentation method (ISO 11277);
- Soil acidity (pH in water extraction 1:5 w/v) – measured potentiometrically (WTW 720 pH meter) (ISO 10390:2002);
- Soil Organic Matter (SOM, %) according to the ISO 14235:1998;
- Total Kjeldahl Nitrogen (TKN, %) content with a modified version of the classic Kjeldahl method (ISO 11261:2002);
- P₂O₅ (mg.100g⁻¹) – extraction with Ammonium Acetate and Calcium Lactate-pH 4.2 (Ivanov, 1984);
- K₂O (mg.100g⁻¹) – extraction with Ammonium Acetate and Calcium Lactate-pH 4.2 (Ivanov, 1984).

The plant species composition was studied. The methods of study included an assessment of the floristic composition, followed by an analysis of the origin and ecological peculiarities of the species. Species identification was carried out according to Delipavlov et al., 2011. Species origin was determined following Walter (Assyov et al., 2012). The experimental plots were of size 4 x 4 m. Floristic affinity among the sites studied was calculated using Sørensen-Dice coefficient (Sørensen, 1948).

The box-plot graphics were created with SPSS 26 (SPSS for MacOS). The linear regressions were done with SPSS 26 (SPSS for MacOS).

The soil parameters were analysed by Analysis of Variance (ANOVA) and Post Hoc LSD ($p \leq 0.05$) test using SPSS 26 (SPSS for MacOS). Different letters are significantly different with level of significance ($p \leq 0.05$).

RESULTS AND DISCUSSION

Plant characteristics. The total number of plant species recorded on the studied sites is 40 as presented in Table 1.

The plant species in our study or research belong to 36 genera and 11 families. The most numerous families are *Poaceae* (30% of species), *Fabaceae* (25%) and *Asteraceae* (23%). Many families (7) as *Boraginaceae*, *Caryophyllaceae* etc. are represented by a single species (less than 3%).

Table 1. List of plant species on the area of Podgumer quarry

Plant species/Coverage	SP 2	SP 3
	70%	80%
<i>Achillea millefolium</i> L.		+
<i>Acinos arvensis</i> (Lam.) Dandy		+
<i>Agrostis capillaris</i> L.	+	
<i>Aegilops triuncialis</i> L.		10%
<i>Bromus mollis</i> L.		+
<i>Bromus squarrosus</i> L.	+	10%
<i>Bromus tectorum</i> L.	+	
<i>Carlina acanthifolia</i> All.	+	
<i>Centaurea rhenana</i> Boreau	+	
<i>Cichorium intybus</i> L.	+	+
<i>Cirsium arvense</i> (L.) Scop.	+	
<i>Coronilla varia</i> L.		+
<i>Cynodon dactylon</i> (L.) Pers.	+	
<i>Dorycnium herbaceum</i> Vill.	+	
<i>Echium vulgare</i> L.	+	+
<i>Equisetum arvense</i> L.	+	
<i>Erysimum diffusum</i> Ehrh.	+	
<i>Euphorbia niciciana</i> Borb.	+	+
<i>Festuca valesiaca</i> Scheich ex Gaud.		20%
<i>Filago vulgaris</i> Lam.	+	
<i>Holcus lanatus</i> L.	+	
<i>Lolium perenne</i> L.		+
<i>Lotus corniculatus</i> L.	1%	+
<i>Medicago lupulina</i> L.	30%	5%
<i>Melilotus alba</i> Medic.		+
<i>Petrorhagia prolifera</i> (L.) P. W. Ball. et Hey.	+	
<i>Poa annua</i> L.		+
<i>Ononis arvensis</i> L.	5%	
<i>Rumex acetosella</i> L.	+	+
<i>Sanguisorba minor</i> Scop.		+
<i>Taeniatherum asperum</i> Nevski		10%
<i>Dasypyrum villosum</i> (L.) Cand.		10%
<i>Thlaspi arvense</i> L.	+	+
<i>Tragopogon dubius</i> Scop.	+	
<i>Trifolium repens</i> L.	10%	10%
<i>Trifolium strictum</i> L.		+
<i>Tussilago farfara</i> L.	20%	
<i>Vicia cracca</i> L.	+	
<i>Vicia sativa</i> L.	+	
<i>Xeranthemum annuum</i> L.	+	

At SP 3 projective coverage is 80% and 22 plant species are observed. SP 2 has a 70% projective coverage and 27 plant species.

Experimental plots differ substantially from each other - Sørensen (1948) coefficient of similarity between plots is 0.33.


The vegetation on embankments goes through different stages of recovery. The formed vegetation on the embankments may be assigned to one of the following stages: 1-weed-ruderal; 2-cereal-ephemeral; 3 – rhizome-leguminous.

At SP 2, most of the observed species are characteristic of the rhizome-leguminous stage. These are species such as: *Holcus lanatus* L., *Cynodon dactylon* (L.) Pers. of cereals and *Medicago lupulina* L., *Lotus corniculatus* L. of legumes. The participation of species from other stages is quite limited.

At SP 3 the predominant species are of rhizome-leguminous stage such as *Trifolium repens* L., *Melilotus alba* Medic. of the legumes and *Festuca valesiaca* Scheich ex Gaud., *Lolium perenne* L. of the rhizome, but at the same time species of the cereal-ephemeral stage are observed, such as *Taeniatherum asperum* Nevski, *Dasypyrum villosum* (L.) Cand.

The analysis of the ratio of biological plant types shows the prevalence of perennial herbaceous plants (55-74%), followed by the group of annual herbaceous plants (36-22%).

There is a normal reduction in the number of annual herbaceous plants at SP 3 to SP 2 since most of them are ruderal species characteristics of the early stages of plant cover formation. Shrubs and tree species are not observed.

**Figure 1.** Share (%) distribution of species biological type

The analysis of the phytogeographical structure of the studied plants reveals that 13 floristic elements are present (Fig. 2). Distribution of species according to their phytogeographic origin shows as most numerous the European geoelements – 25% of species. Other groups with larger participation (about 13% each) are: Eur-Sib, Eur-Med, subMed. There are 5 groups represented by one species (around 3%), such as: EursubMed, PontMed, Pont, Med and CSEur. Adventitious species were not observed and other groups with greater participation - about 13% each are: Eur-Sib, Eur-Med, subMed.

Soil characteristics

There are statistically significant differences in the bulk densities of the materials deposited in the different years, where the change is expressed as a statistically significant correlation (Fig. 3). Total porosity increases with time and the change is statistically significant (Fig. 4).




Figure 2. Share (%) ratio distribution of floristic elements




Figure 3. Box-plot graph and a linear regression of BD depending on the year of deposit




Figure 4. Box-plot graph and a linear regression of TP depending on the year of deposit

There are statistically significant differences in the three fractions of the soil texture in the different years of deposit. The decrease in the sand fraction (Fig. 5), results in the increase in the silt and clay fractions, where the silt fraction has a greater rate of change than the clay fraction. Similar trends (decrease in the sand fraction, increase in the silt fraction and lesser rate of change in the clay fraction) were reported by Bogdanov (2014b). Nevertheless, soil texture is considered to be stable and can change very slowly with time (Atanasov, 1987).




Figure 5. Soil texture changes (USDA 2017)

There are no statistically significant differences in the active soil acidity between the year of deposit and 5 years after deposit, but there is a statistically significant difference 9 years after deposit (Fig. 6).

There are statistically significant differences in SOM and TKN in different years of deposit (Fig. 7, Fig. 8), where the changes in the two indicators are expressed by statistically significant linear relationships. The positive correlation between the ages of the site for these two indicators has been discussed by other authors (Frouz et al., 2008).

The plant-available P and K increase with every following year of deposit, where these changes in both indicators are expressed by statistically significant linear correlations (Fig. 9, Fig. 10), which is consistent with studies done by other authors (Frouz et al., 2008), who report statistically significant positive correlations between the age of the site and forms of P and K.




Figure 6. Box-plot graph of pH in the different years of deposit




Figure 7. Box-plot graph and a linear regression of SOM depending on the year of deposit




Figure 10. Box-plot graph and a linear regression of K₂O depending on the year of deposit




Figure 8. Box-plot graph and a linear regression of TKN depending on the year of deposit




Figure 9. Box-plot graph and a linear regression of P₂O₅ depending on the year of deposit

The natural restoration of a site is a long process and it is sometimes advisable to study the dynamics for a certain period of time before we adopt measures for the management of post-mining areas (Letheren, 2009; Rao et al., 2000). The changes in plant cover (pioneer species to begin with) are consistent with the observations made by other authors (Harris et al., 1998; Tomlinson et al., 2008; Walker and del Moral, 2003). However, spontaneous succession is difficult to predict in the context of global change. For that reason, it is not considered to be an acceptable rehabilitation strategy for post-mining areas (Cullen et al., 1998). The rehabilitation of ecosystems and ecological functions in a sandy landscape requires soil reconstruction (Jim, 2001) and therefore landscape planning measures should be taken (Bradshaw, 1997).

CONCLUSIONS

The establishment of rhizome-leguminous plants and the changes in physical and chemical properties point to the presence of an early soil formation process. The vegetation had a significant impact (statistically significant) on bulk density (decrease) 5 years and 9 years after deposit (described by reliable linear regressions). There was a statistically significant increase in total porosity after the establishment of vegetation (described by reliable linear regressions).

The establishment of vegetation resulted in statistically significant differences in the chemical characteristics (SOM, TKN, P₂O₅, K₂O). The establishment of vegetation and the increase in plant diversity resulted in increased levels of SOM, TKN, P₂O₅, K₂O, where these changes were described by reliable linear regressions.

ACKNOWLEDGEMENT

The authors would like to extend their sincere gratitude to the National Research Programme “Young Scientists and Post-doctoral Candidates” for their support in the implementation of parts of this study.

REFERENCES

- Atanasov I. (1978). *Soil Science and Tropical Soils*. Zemizdat, Sofia, p. 260.
- Assyov B., Petrova A. (2012). *Conspectus of the Bulgarian Vascular Flora. Horology and Floristic Elements*. Bulgarians Biodiversity Foundation, Sofia, p. 492.
- Berendse F., Lammerts E., Olff H. (1998). Soil Organic Matter Accumulation and Its Implications for Nitrogen Mineralization and Plant Species Composition during Succession in Coastal Dune Slacks. *Plant Ecol.*, 137: 71 - 78. DOI: doi.org/10.1023/A:1008051931963
- Bogdanov S. (2013). Changes in the Acidity of Luvisols under the Influence of Forest Fires, *Ecological Engineering and Environmental Protection*, 2: 61-67.
- Bogdanov S. (2014a). Plant Available Water Capacity as a Factor for the Sustainable Development of Forests Affected by Forest Fires. *Management and Sustainable Development*, 3(46): 23-26.
- Bogdanov S. (2014b). Soil Texture Changes in Gray Forest Soils (Gray Luvisols) Influenced by Forest Fires in Deciduous Forests. *Forestry Ideas*, Vol. 20, No 2 (48): 135-140
- Bowen C.K., Schuman G., Olson R., Ingram J. (2005). Influence of Topsoil Depth on Plant and Soil Attributes of 24-year Old Reclaimed Mined Lands. *Arid Land Res. Manag.*, 267-284; DOI: dx.doi.org/10.1080/15324980590951441.
- Bradshaw A. (1997). Restoration of Mined Lands, Using Natural Processes. *Ecol. Eng.* (8): 255-269; DOI: dx.doi.org/10.1016/S0925-8574(97)00022-0.
- Bradshaw A. (2000). The Use of Natural Processes in Reclamation – Advantages and Difficulties. *Landscape and Urban Planning*, 51: 89-100; DOI: 10.1016/S0169- 2046(00)00099-2.
- Bradshaw A.D., Huttl R.F. (2001). Future Minesite Restoration Involves a Broader Approach. –*Ecol. Eng.* 17: 87-90; DOI: dx.doi.org/10.1016/S0925-8574(00)00149-X.
- Chenot, J., Jaunatre, R., Buisson, E., Bureau, F., Dutoit, T. (2018). Impact of Quarry Exploitation and Disuse on Pedogenesis. *Catena*, 160, 354-365. DOI: doi.org/10.1016/j.catena.2017.09.012
- Corry R.C., Laforteza R., Brown R.D. (2010). Ecological Functionality of Landscapes with Alternative Rehabilitations of Depleted Aggregate Sites. *Mining Reclam Environ.* 24: 216-232.
- Darwish T., Khater C., Jomaa I., Stehouwer R., Shaban A., Hamze M. (2011). Environmental Impact of Quarries on Natural Resources in Lebanon. *Land Degrad.* 22: 345–358; DOI: dx.doi.org/10.1002/ldr.1011.
- De Deyn G.B., Raaijmakers C.E., Zoomer H.R., Berg M.P., Ruiter P.C., Verhoef A., Bezemer T.M., Putten W.H. (2003). Soil Invertebrate Fauna Enhances Grass-Land Succession and Diversity. *Nature*, 422: 711-713; DOI: dx.doi.org/10.1038/nature01548.
- Delipavlov D., Cheshmedjiev I., Popova M., Terzijski D., Kovachev I. (2011). *Guide of Vascular Plants in Bulgaria*. Agronomy University, Plovdiv, p. 591.
- Deng X., Liu Y., Han S. (2009). Carbon and Nitrogen Dynamics in Early Stages of Forest Litter Decomposition as Affected by Nitrogen Addition. *J. Forest Res.*, 20: 111-116. DOI: doi.org/10.1007/s11676-009-0020-4
- Frouz J., Prach K., Pižl V., Háněl L., Starý J., Tajovský K., Materna J., Balik V., Kalcik J., Řehounková K. (2008). Interactions between Soil Development, Vegetation and Soil Fauna during Spontaneous Succession in Post Mining Sites. *European Journal of Soil Biology*, 44(1): 109–121. doi:10.1016/j.ejsobi.2007.09.002
- Geissen V., Wang S., Oostindie K., Huerta E., Zwart K., Smit A., Ritsema C., Moore D. (2013). Effects of Topsoil Removal as a Nature Management Technique on Soilfunctions. *Catena*, 101: 50-55; DOI: dx.doi.org/10.1016/j.catena.2012.10.002.
- Harris J.A., Birch P., Palmer J.P. (1998). *Land Restoration and Reclamation: Principles and Practice*. Addison Wesley Longman, Harlow.
- Hristov, B. (2020). The Soils of Botevgrad Valley. *Ecol. Engin. and Env. Protect.*, (2): 52-62
- Hristova M. (2013). *Content and Availability of Microelements-Metals in Technogenic Soils*. Poushkarov, Sofia, p. 143
- Heneghan L., Miller S., Baer S., Callahan M., Montgomery J., Pavao-Zuckerman M., Rhoades C., Richardson S. (2008). Integrating Soil Ecological Knowledge into Restoration Management. – *Restor. Ecol.* (16): 608-617; DOI: dx.doi.org/10.1111/j.1526-100X.2008.00477.x.
- Holl K.D. (2002). Long-term Vegetation Recovery on Reclaimed Coal Surface Mines in The Eastern USA. *Journal of Applied Ecology*, 39: 960-970; DOI: 10.1046/j.1365- 2664.2002.00767.x.
- ISO. (1998). *Soil Quality. Determination of Organic Carbon by Sulphochromic Oxidation*. International Organization for Standardization (ISO 14235). p. 5. Geneva, Switzerland
- ISO. (2001). *Soil Quality. Determination of Dry Bulk Density*. International Organization for Standardization (ISO 11272). p. 14 Geneva, Switzerland
- ISO. (2002). *Soil Quality. Determination of Particle Size Distribution in Mineral Soil Material, Method by Sieving and Sedimentation*. International Organization for Standardization (ISO 11277). p. 10. Geneva, Switzerland
- ISO. (2002). *Soil Quality. Determination of pH*. International Organization for Standardization (ISO 10390). p. 8. Geneva, Switzerland
- ISO. (2002). *Soil Quality. Determination of Total Nitrogen - Modified Kjeldahl Method*. International Organization for Standardization (ISO 11261). p. 7. Geneva, Switzerland
- IUSS. (2014). *World Reference Base for Soil Resources. International Soil Classification System for Naming Soils and Creating Legends for Soil Maps*. World Soil Resources Reports No. 106. Working Group WRB. FAO, Rome.
- Ivanov P. (1984). New Method to Determine the Plant Available Phosphorus and Potassium in Soil. – *Soil and Agrochemistry*, 4: 88-98.
- Jian G. Y., Fang W., Fan L., Chen Y., Wang D., Yang Z. (2006). Soil Formation and Vegetation Establishment on the Cliff Face of Abandoned Quarries in the Early Stages of Natural Colonization. – *Restoration Ecology Volume 14*(3): 349- 356. DOI: doi.org/10.1111/j.1526-100X.2006.00143.x
- Krummelbein J., Bens O., Raab T., Naeth A. 2012. A History of Lignite Coal Mining Reclamation Practices in Lusatia, Eastern Germany. *Canadian Journal of Soil Science*, 92: 55–66; doi: 10.4141/cjss2010-063.
- Larson E., O'Malley D., Melloni R. (2006). Aggression and Vasotocinane Associated with Dominant-Subordinate Relationships in Zebrafish. *Behavioural Brain Research*. 167(1): 94-102.
- Lorraine F., Flint A. (2002). Porosity. - In: Dane J., G. Topp (Eds.) *Methods of Soil Analysis, Part 4. Physical methods*. No. 5. Madison, WI, Soil Science Society of America, 241-253.
- Mouflis G.D., Gitas I.Z., Iliadou S., Mitri G.H. (2008). Assessment of the Visual Impact of Marble Quarry Expansion (1984–2000) on the Landscape of Thasos Island, NE Greece. *Landsc. Urban Plan.*, 86: 92-102; DOI: dx.doi.org/10.1016/j.landurbplan.2007.12.009.
- Nieman T., Merkin Z. (1995). *Wildlife Management, Surface Mining, and Regional Planning*. – *Growth and Change*, 26(3): 405-424. DOI: doi.org/10.1111/j.1468-2257.1995.tb00178.x
- Petersen R.G., Calvin L.D. (1996). *Sampling*. – In: J. M. Birgham (Ed.) *Methods of Soil Analysis, Part 3. Chemical Methods*, No. 5. Madison, WI, Soil Science Society of America, 1-19.
- Pietrzykowski M., Krzaklewski W. (2007). An Assessment of Energy Efficiency in Reclamation to Forest. – *Ecol. Eng.* 30: 341 - 348. DOI: doi.org/10.1016/j.ecoleng.2007.04.003
- Prach K., Řehounková K., Řehounek J., Konvalinková P. (2011). Ecological Restoration of Central European Mining Sites: a Summary of a Multi-Site Analysis. *Landscape Research*, 36: 263-268; DOI: 10.1080/01426397.2010.547571.
- Prieto B., Silva B., Aira N., Laiz L. (2005). Induction of Biofilms on Quartz Surfaces as a Means of Reducing the Visual Impact of Quartz Quarries. *Biofouling*, 21(5-6): 237-246; DOI: 10.1080/08927010500421294.

- Prieto B., Silva B., Aira N. (2006). Methodological Aspects of the Induction of Biofilms for Remediation of the Visual Impact Generated by Quartz Mining. *Science of the Total Environment*, 370 (1): 254 -261. DOI: doi.org/10.1016/j.scitotenv.2006.07.030
- Primack R., Sarrazin B., Lecomte F. (2012). *Biologie de la Conservation*. Dunod, Paris, France, p. 360.
- Romana M. (1985). New Adjustment Ratings for Application of Bieniawski Classification to Slopes. *Proceedings of the International Symposium on the Role of Rock Mechanics*. – Zacatecas, Mexico, 49-53.
- Schulz D. (1996). Recultivation of Mining Waste Dumps in the Ruhr Area, Germany. *Water Air Soil Pollut*, 91: 89-98. DOI: doi.org/10.1007/978-94-015-8694-8_7
- Sebelikova L., Csicssek G., Kirmer A., Vitovcova K., Ortmann-Ajkai A., Prach K., RŘehounek K. (2018). Spontaneous Revegetation vs. Forestry Reclamation - Vegetation Development in Coal Mining Spoil Heaps across Central Europe. *Land Degradation & Development*; 30(3), 348-356; DOI:10.1002/ldr.3233; 30(3): 348-356.
- Singh A., Raghubanshi A., Singh J. (2004). Comparative Performance and Restoration Potential of Two Albizia Species Planted on Mine Spoil in a Dry Tropical Region, India. *Ecological Engineering*. 22(2): 123-140. DOI: doi.org/10.1016/j.ecoleng.2004.04.001
- Sorensen T. (1948). A Method of Establishing Groups of Equal Amplitude in Plant Sociology Based on Similarity of Species and Its Application to Analyses of the Vegetation on Danish Commons. *Kongelige Danske Videnskabernes Selskab*, 5(4): 1-34.
- Sourkova M., Frouz J., Santruckova H. (2005). Accumulation of Carbon, Nitrogen and Phosphorus During Soil Formation on Alder Spoil Heaps After Brown-Coal Mining Near Sokolov (Czech Republic). *Geoderma*, 124: 203-214. DOI: doi.org/10.1016/j.geoderma.2004.05.001
- Stahl P., Perryman D., Sharmasarkar B., Munn S. (2002). Topsoil Stock Piling versus Exposure to Traffic: A Case Study on in Situ Uranium Wellfields. *Restor. Ecol.* 10, 129-137; DOI: dx.doi.org/10.1046/j.1526-100X.2002.10114.x.
- Torreta N., Takeda H. (1999). Carbon and Nitrogen Dynamics of Decomposing Leaf Litter in a Tropical Hill Evergreen Forest. *Eur. J. Soil Biol.*, 35: 57-63. DOI: doi.org/10.1016/S1164-5563(99)00111-9
- Van-Camp L., Bujarrabal B., Gentile A. R., Jones R. J. A., Montanarella L., Olazabal C., Selvaradjou S. (2004). Reports of the Technical Working Groups Established Under the Thematic Strategy for Soil Protection. EUR 21319 EN/1. Office for Official Publications of the European Communities, Luxembourg. p. 79.
- Walker L. R., Moral R. (2003). *Primary Succession and Ecosystem Rehabilitation*. Cambridge University, p. 426.
- Zahariev B. (1979). *Forest Vegetation Zoning of the People's Republic of Bulgaria*. Zemizdat. p. 195.
- Zhuang X.Y. (1997). Rehabilitation and Development of Forest on Degraded Hills of Hong Kong. *Forest Ecol. Manage*, 99:197 - 201. DOI: doi.org/10.1016/S0378-1127(97)00205-3
- USDA. (2017). *Soil Survey Manual*. Soil Science Division Staff. Agriculture Handbook No. 18.

aCS86_2